DISEASE:
Hyperinsulinism due to HNF1A deficiency

NAME:	Hyperinsulinism due to HNF1A deficiency
DESCRIPTION:	Hyperinsulinism due to HNF1A deficiency is a form of diazoxide-sensitive diffuse hyperinsulinism (DHI), characterized by transient or persistent hyperinsulinemic hypoglycemia (HH) in infancy that is responsive to diazoxide, evolving in to maturity-onset diabetes of the young subtype 1 (MODY-1; see this term) later in life.
ORPHACODE:	324575
SYNONYMS:	Hyperinsulinemic hypoglycemia due to HNF1A deficiency
XREF(S):	Orphanet
ANALYTE(S):	HNF1A
CREATED:	13 May 2019 - 01:02
CHANGED:	22 Jun 2023 - 16:14

RELATED CONTENT

Related Genetic Tests

- Diabetes neonatal / Maturity onset Diabete of the Young (MODY) / Hyperinsulinism (gene panel)
- Hyperinsulinism (gene panel)
- Maturity onset Diabete of the Young (MODY), type 5 / Renal cysts and diabetes syndrome (gene panel)

Related Laboratories

- Centrum Medische Genetica - UZ Antwerpen

Related Analytes

- HNF1 homeobox A

Related Gene Panels

- Diabetes neonatal / Maturity onset Diabete of the Young (MODY) / Hyperinsulinism (genepanel) - UZA
- MODY (7 genes) - UZA