

DISEASE:
Pituitary stalk interruption syndrome

NAME:	Pituitary stalk interruption syndrome
DESCRIPTION:	Pituitary stalk interruption syndrome (PSIS) is a congenital abnormality of the pituitary that is responsible for pituitary deficiency and is usually characterized by the triad of a very thin or interrupted pituitary stalk, an ectopic (or absent) posterior pituitary (EPP) and hypoplasia or aplasia of the anterior pituitary visible on MRI. In some patients the abnormality may be limited to EPP (also called ectopic neurohypophysis) or to an interrupted pituitary stalk.
ORPHACODE:	95496
SYNONYMS:	Ectopic neurohypophysis PSIS
XREF(S):	<u>Orphanet</u> <u>ICD-10</u>
ANALYTE(S):	<u>WDR11</u> <u>ROBO1</u> <u>PROKR2</u> <u>HESX1</u> <u>LHX4</u> <u>CDON</u> <u>GPR161</u>

CREATED:	13 May 2019 - 01:02
CHANGED:	22 Jun 2023 - 16:14

Source URL: <http://gentest.healthdata.be/disease/3444>

RELATED CONTENT

Related Genetic Tests

- [Brain malformations \(gene panel\)](#)
- [Hypogonadotropic hypogonadism \(33 genes\)](#)

Related Laboratories

- [Centre de Génétique Humaine - Erasme ULB](#)
- [Centrum Medische Genetica - UZ Brussel VUB](#)

Related Analytes

- [cell adhesion associated, oncogene regulated](#)
- [G protein-coupled receptor 161](#)
- [HESX homeobox 1](#)
- [LIM homeobox 4](#)
- [prokineticin receptor 2](#)
- [roundabout guidance receptor 1](#)
- [WD repeat domain 11](#)

Related Gene Panels

- Hypogonadotropic Hypogonadism/Kallmann (61 genes) - ULG
- Hypogonadotropic hypogonadism (33 genes) - VUB

Source URL: <http://gentest.healthdata.be/disease/3444>