

GENETIC TEST:
Thrombophilia due to protein C deficiency (PROC gene)

FULL NAME:	Thrombophilia due to protein C deficiency (PROC gene)
TEST TYPE:	Clinical
TEST SPECIALTY:	Molecular Genetics
TEST PURPOSE:	Carrier diagnosis, Mutation confirmation, Post-natal Diagnosis, Prenatal diagnosis
SPECIMEN:	Peripheral (whole) blood on EDTA, Amniotic fluid, Chorionic villi, Cell culture
METHOD CATEGORY:	Sequence analysis: entire coding region Mutation screening and sequence analysis of selected exons
METHOD TECHNIQUE:	Bi-directional Sanger Sequence analysis
RIZIV CODE:	565456-565460

EQA:	• DNA Sequencing - Sanger, • DNA Sequencing - Sanger, • DNA Sequencing - Sanger, • DNA Sequencing - Sanger, • DNA Sequencing - Sanger, • DNA Sequencing - Sanger, • DNA Sequencing - Sanger , • DNA Sequencing - Sanger
TURNAROUND TIME (MAXIMUM):	3 months (10 working days for prenatal diagnosis)
CREATED:	28 Aug 2019 - 14:18
CHANGED:	09 Mar 2023 - 16:26
URL:	https://laboguide.uzbrussel.be/laboguide#Analyses:Protein%20C%20defici%C3%ABnti...

Source URL: http://gentest.healthdata.be/genetic_test/790

RELATED CONTENT

Related Diseases

- [Severe hereditary thrombophilia due to congenital protein C deficiency](#)

Related Laboratories

- [Centrum Medische Genetica - UZ Brussel VUB](#)

Related Analytes

- [protein C, inactivator of coagulation factors Va and VIIIa](#)

Source URL: http://gentest.healthdata.be/genetic_test/790